arXiv:0801.0495v2 [math.CO] 31 Jul 2010

TORIC IDEALS OF FLOW POLYTOPES

MATTHIAS LENZ

Abstract. We show that toric ideals of flow polytopes are generated in degree 3. This was conjectured by Diaconis and Eriksson for the special case of the Birkhoff polytope. Our proof uses a hyperplane subdivision method developed by Haase and Paffenholz.

It is known that reduced revlex Gröbner bases of the toric ideal of the Birkhoff polytope B_n have at most degree n. We show that this bound is sharp for some revlex term orders. For $(m \times n)$-transportation polytopes, a similar result holds: they have Gröbner bases of at most degree $\lfloor mn/2 \rfloor$. We construct a family of examples, where this bound is sharp.

1. Introduction

Let $G = (V, A)$ be a directed graph and $\boldsymbol{d} \in \mathbb{Z}^V$, $\boldsymbol{l}, \boldsymbol{u} \in \mathbb{N}^A$. A *flow* on G is a function $f : A \to \mathbb{R}_{\geq 0}$ that respects the lower and upper bounds $\boldsymbol{l}$ and $\boldsymbol{u}$ and satisfies the demand $\boldsymbol{d}$, *i. e.* for every vertex v, the flow entering v minus the flow leaving v equals d_v. A *flow polytope* is the set of all flows with fixed parameters $G, \boldsymbol{d}, \boldsymbol{u}, \boldsymbol{l}$.

An important special case are *transportation polytopes*. They can be written as sets of $(m \times n)$-matrices whose row and column sums equal some fixed positive integers.

Given a lattice polytope P, the relations among the lattice points in P define the *toric ideal* I_P. Generating sets of toric ideals are used in algebraic statistics to construct Markov chain algorithms for sampling from conditional distributions ([2]). In particular, small generating systems that can be handled by computers are of practical interest. *Diaconis and Eriksson* ([1]) conjectured that the toric ideal of the Birkhoff polytope B_n (the convex hull of all $(n \times n)$-permutation matrices) is generated in degree 3. They proved this by massive computations for $n \leq 6$. For arbitrary $n \geq 4$, they showed that I_{B_n} has a generating set of degree $n - 1$.

Haase and Paffenholz ([5]) proved that the toric ideals of almost all (3×3)-transportation polytopes and the smooth ones in particular are generated in degree 2. The only exception is the Birkhoff polytope B_3, whose toric ideal is generated in degree 3.

Our Main Theorem proves the Diaconis-Eriksson conjecture and generalizes the result of Haase and Paffenholz:

Theorem 1.1 (Main Theorem)**.** *Toric ideals of flow polytopes are generated in degree* 3.

1991 *Mathematics Subject Classification.* Primary 52B20; Secondary 13P10, 14M25.

Key words and phrases. Toric ideal, Flow polytope, Transportation polytope, Gröbner basis, Markov basis.

Toric ideals define toric varieties, which are an important class of examples in algebraic geometry. A lattice polytope P is smooth if the edge directions at every vertex form a lattice basis. This is equivalent to the corresponding toric variety X_P being smooth. It was conjectured that if X_P is smooth, the defining ideal is generated by quadratic binomials or even possesses a quadratic Gröbner basis ([3], [13, Conjecture 2.9]). The original motivation of our research was to check if this conjecture holds for flow polytopes.

In *Section 2*, we review some important definitions and theorems. In *Section 3*, we describe a method that can be used to prove degree bounds for generating sets and Gröbner bases of toric ideals: first, we choose a nice triangulation of the point set and then, we use a correspondence between Gröbner bases and triangulations established by *Sturmfels*. This hyperplane subdivision method was developed by Haase and Paffenholz in [5]. In *Section 4*, we apply the method described in Section 3 to prove our Main Theorem.

Diaconis and Sturmfels showed that all revlex Gröbner bases of I_{B_n} are at most of degree n ([2, Theorem 6.1], [12, Theorem 14.8]). Computational experiments provide evidence that this bound might be optimal ([1, Remark 9]). In *Section 5* we show that for some revlex term orders, this bound is indeed optimal. Both the bound and the examples can be generalized to $(m \times n)$-transportation polytopes: reduced Gröbner bases with respect to a certain class of term orders are at most of degree $\lfloor mn/2 \rfloor$ and we construct a family of transportation polytopes and term orders, where this bound is almost sharp.

2. Background

In this section, we review some important definitions and theorems.

2.1. **Notation.** $\mathbb{N} = \{0, 1, 2, \ldots\}$. Matrices are denoted by capital letters, vectors by boldface small letters. Their entries are denoted by the corresponding small letters. Let $\boldsymbol{a} = (a_1, \ldots, a_n) \in \mathbb{N}^n$. We write $\boldsymbol{x}^{\boldsymbol{a}}$ to denote the monomial $\prod_{i=1}^n x_i^{a_i} \in k[\boldsymbol{x}] = k[x_1, \ldots, x_n]$.

The term *polytope* always refers to a convex lattice polytope, *i. e.* all vertices of our polytopes are integral. For background information on polyhedral geometry and polytopes see *Schrijver*'s or *Ziegler*'s book [11, 15].

2.2. **Flow polytopes.** *Flow polytopes* (or transshipment polytopes) are the main geometric objects we are dealing with. Let $G = (V, A)$ be a directed graph and $\boldsymbol{d} \in \mathbb{Z}^V$, $\boldsymbol{l}, \boldsymbol{u} \in \mathbb{N}^A$. Let $M_G \in \{-1, 0, 1\}^{V \times A}$ denote the vertex-arc incidence matrix of G.

Note that the definition of *flow polytopes* given in the introduction is equivalent to

$$F = F_G = F_{G,\boldsymbol{d},\boldsymbol{u},\boldsymbol{l}} = \left\{ \boldsymbol{f} \in \mathbb{R}^A_{\geq 0} \,\middle|\, M_G \boldsymbol{f} = \boldsymbol{d}, \, \boldsymbol{l} \leq \boldsymbol{f} \leq \boldsymbol{u} \right\} \tag{1}$$

It is a standard fact that M_G is totally-unimodular. This implies that the polytope F has integral vertices.

Throughout this paper, we suppose that all our flow polytopes F are *homogeneous*, *i. e.* F is contained in an affine hyperplane, that does not contain the origin. If $\boldsymbol{d} \neq \boldsymbol{0}$ this statement holds. Otherwise, we consider the homogenized polytope $\{1\} \times F$.

An important special case of flow polytopes are transportation polytopes. In statistics, they appear under the name 2*-way contingency tables.*

Let $m, n \in \mathbb{Z}_{\geq 1}, \boldsymbol{r} \in \mathbb{Z}_{\geq 1}^m, \boldsymbol{c} \in \mathbb{Z}_{\geq 1}^n$ be two vectors satisfying $s := \sum_{i=1}^n c_i = \sum_{i=1}^m r_i$. The *transportation polytope* $T_{\boldsymbol{rc}}$ is defined as

$$T_{\boldsymbol{rc}} = \left\{ A \in \mathbb{R}_{\geq 0}^{m \times n} \,\middle|\, \sum_{i=1}^m a_{ij} = c_j, \sum_{j=1}^n a_{ij} = r_i \right\} \tag{2}$$

The upper $((m-1) \times (n-1))$-minor of a matrix $A \in T_{\boldsymbol{rc}}$ determines all other entries. Hence, the dimension of $T_{\boldsymbol{rc}}$ is at most $(m-1)(n-1)$. On the other hand, $a_{ij} = r_i c_j / s$ determines an interior point, so that the dimension is exactly $(m-1)(n-1)$.

If $\boldsymbol{r} = \boldsymbol{c} = (1, \ldots, 1)$, we obtain an important example: the Birkhoff polytope B_n. We will use the following theorem in the proof of the Main Theorem:

Theorem 2.1 (Generalized Birkhoff-von Neumann Theorem)**.**
Let $G = (V, A)$ be a directed graph and let $F = F_{G,\boldsymbol{d},\boldsymbol{u},\boldsymbol{l}}$ be a flow polytope and $k \in \mathbb{N}$. Let $\boldsymbol{f} \in (k \cdot F) \cap \mathbb{Z}^A$. Then, there exist flows $\boldsymbol{f}_1, \ldots, \boldsymbol{f}_k \in F \cap \mathbb{Z}^A$ s.t. $\boldsymbol{f} = \boldsymbol{f}_1 + \ldots + \boldsymbol{f}_k$.

For a combinatorial proof, see *Schrijver* [11, Th. 13.14]. Readers that are familiar with lattice polytopes might prefer the following proof: flow polytopes admit a unimodular triangulation (see below). Polytopes that admit a unimodular triangulation are normal.

2.3. **Toric ideals and Gröbner bases.** This paragraph defines toric ideals and Gröbner bases as in *Sturmfels*'s book ([12]). Let k be a field and let $P \subseteq \mathbb{R}^d$ be a homogeneous lattice polytope. The set of its lattice points $\mathcal{A}$ defines a semigroup homomorphism $\pi : \mathbb{N}^{\mathcal{A}} \to \mathbb{Z}^d, \boldsymbol{u} \mapsto \sum_{\boldsymbol{a} \in \mathcal{A}} u_{\boldsymbol{a}} \boldsymbol{a}$, which can be lifted to a ring homomorphism

$$\hat{\pi} : k[x_{\boldsymbol{a}} \mid \boldsymbol{a} \in \mathcal{A}] \to k[t_1^{\pm 1}, \ldots t_n^{\pm 1}], \qquad x_{\boldsymbol{a}} \mapsto \mathbf{t}^{\boldsymbol{a}} \tag{3}$$

Its kernel is the homogeneous ideal $I_{\mathcal{A}} = \left\langle \mathbf{x}^{\mathbf{u}} - \mathbf{x}^{\mathbf{v}} \;\middle|\; \sum_{\boldsymbol{a} \in \mathcal{A}} u_{\boldsymbol{a}} \boldsymbol{a} = \sum_{\boldsymbol{a} \in \mathcal{A}} v_{\boldsymbol{a}} \boldsymbol{a} \right\rangle$. This ideal is called the *toric ideal* associated to $\mathcal{A}$ (or P respectively).

A binomial $\boldsymbol{x}^{\boldsymbol{u}} - \boldsymbol{x}^{\boldsymbol{v}} \in I_{\mathcal{A}}$ corresponds to a relation between points in $\mathcal{A}$. For example, for $\mathcal{A} = B_3 \cap \mathbb{Z}^{3 \times 3}$, $I_{\mathcal{A}}$ is generated by the binomial that corresponds to the relation $\sum_{\det(M)=1} M = \sum_{\det(M)=-1} M$.

A total order $\prec$ on $\mathbb{N}^n$ is a *term order* if $\boldsymbol{a} \prec \boldsymbol{b}$ implies $\boldsymbol{a} + \boldsymbol{c} \prec \boldsymbol{b} + \boldsymbol{c}$ for all $\boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c} \in \mathbb{N}^n$ and the zero vector is the unique minimal element. An important example is the graded reverse lexicographic (revlex) order: $\boldsymbol{a} \prec_{\text{revlex}} \boldsymbol{b}$ if $\sum_i a_i < \sum_i b_i$ or $\sum_i a_i = \sum_i b_i$ and the rightmost non-zero entry in $\boldsymbol{a} - \boldsymbol{b}$ is positive. Note that the revlex order depends on the order of the variables.

Let $I \subseteq k[x_1, \ldots, x_n]$ be an ideal and let $\prec$ denote a term order on $\mathbb{N}^n$. For $f \in k[x_1, \ldots, x_n]$, let $\text{in}_\prec(f)$ denote the initial (largest) term of f with respect to $\prec$. A finite set $\mathcal{G} \subseteq I$ is a Gröbner basis of I if for every $f \in I$, there exists a $g \in \mathcal{G}$ s. t. $\text{in}_\prec(g) | \text{in}_\prec(f)$. A Gröbner basis $\mathcal{G}$ is called *reduced* if for two distinct elements $g, g' \in \mathcal{G}$, no term of g' is divisible by $\text{in}_\prec(g)$. The reduced Gröbner basis is unique if we fix an ideal and a term order and require the coefficient of the initial term of every element to be 1.

For $\omega \geq 0$ and an arbitrary term order $\prec$, we obtain a new term order $\prec_\omega$ by $\boldsymbol{a} \prec_\omega \boldsymbol{b}$ if and only if $\boldsymbol{a} \cdot \omega < \boldsymbol{b} \cdot \omega$ or ($\boldsymbol{a} \cdot \omega = \boldsymbol{b} \cdot \omega$ and $\boldsymbol{x}^{\boldsymbol{a}} \prec \boldsymbol{x}^{\boldsymbol{b}}$).

2.4. Regular triangulations and initial ideals.

We assume that the reader is familiar with *regular* and *pulling triangulations* (see *e. g.* [6] or [9]). A triangulation is called *unimodular* if the normalized volume of all simplices contained in it equals 1.

For a generic $\omega \in \mathbb{R}^n$, there is a connection between the regular triangulation and the term order defined by ω. This is described by the following theorem, which is a conglomerate of Corollaries 8.4 and 8.9 in Sturmfels's book ([12]):

Theorem 2.2. *Let P be a polytope and Δ be a regular, unimodular triangulation of P. Let $\prec_\Delta$ be the term order corresponding to Δ. Then, the initial ideal of I_P with respect to $\prec_\Delta$ is given by*

$$\operatorname{in}_{\prec_\Delta}(I_P) = \left\langle \boldsymbol{x}^F \mid F \textit{ is a minimal non-face of } \Delta \right\rangle \tag{4}$$

Let Δ be a regular triangulation that is induced by the vector ω. It is known that Δ is a pulling triangulation if and only if $\prec_\omega$ is the reverse lexicographic term order.

3. The Hyperplane Subdivision Method

In this section, we describe a general method for showing degree bounds for generating sets and Gröbner bases of toric ideals that was developed by Haase and Paffenholz ([5]).

A flow polytope has a canonical subdivision into polytopes contained in lattice translates of a unit cube: we slice F along hyperplanes of type $H_{ak} = \{\boldsymbol{x} \mid x_a = k\}$. Let $F = F_{G,\boldsymbol{d},\boldsymbol{u},\boldsymbol{l}}$ be a flow polytope. For $\boldsymbol{k} \in \mathbb{Z}^A$ we define a *cell of* F as $Z_F(\boldsymbol{k}) = \{\boldsymbol{f} \in F \mid k_a \le f_a \le k_a + 1 \text{ for all } a \in A\}$. Cells are flow polytopes using the same graph with tighter upper and lower bounds and thus lattice polytopes. For our purposes, it is acceptable to identify a cell with the translated cell $Z_F(\boldsymbol{k}) - \boldsymbol{k} \subseteq [0,1]^A$.

We will use a particular class of regular triangulations that we call *subdivide-and-pull triangulations*. They are obtained in the following way: start by subdividing the flow polytope along hyperplanes into cells as defined above. Then refine to a triangulation by pulling all vertices.

The following lemma will be used in the next section to prove the Main Theorem:

Lemma 3.1. Let F be a flow polytope and $k \ge 2$. I_F is generated in degree k if the toric ideal of every cell of F is generated in degree k.

Let Δ be a subdivide-and-pull triangulation of F and let $\mathcal{G}$ be the reduced Gröbner basis with respect to the term order $\prec_\Delta$. $\mathcal{G}$ has at most degree k if for every cell Z, the Gröbner basis of I_Z with respect to the restriction of $\prec_\Delta$ to Z has at most degree k.

Proof of Lemma 3.1. Let Δ_F be a subdivide-and-pull triangulation of F. A polytope is said to have facet width 1 if for each of its facets, the polytope lies between the hyperplane spanned by this facet and the next parallel lattice hyperplane. It is known, that a polytope has facet width 1 if and only if all its pulling triangulations are unimodular ([8, 10, 14]]). Cells of flow polynomials have facet width 1. Hence, Δ_F is unimodular.

Observe that all minimal non-faces that are *not* contained in a cell have cardinality 2 ([4]). Both statements of the theorem follow by applying Theorem 2.2. □

Example 3.2. Let $\mathcal{A} = \{a, b, c, d, e, f\} \subseteq \mathbb{Z}^2$ be the set of lattice points in Figure 1. The subdivide-and-pull triangulation in the picture is induced by the height vector $\omega = (9, 0, 10, 9, 0, 10)$. The Gröbner basis with respect to $\prec_\omega$ is $\mathcal{G} = \{\underline{bf} - ae, \underline{ce} - bd, \underline{cf} - be, \underline{ad} - be, \underline{ac} - b^2, \underline{df} - e^2\}$. The initial terms are underlined.



Figure 1. A regular triangulation

4. Proof of the Main Theorem

In this section, we prove that toric ideals of flow polytopes are generated in degree 3 by applying the method described in the previous section. First, we prove our result for transportation polytopes. In the second subsection, we show that the general statement can be reduced to this special case.

4.1. **Proof for Transportation Polytopes.**

Theorem 4.1. *Toric ideals of transportation polytopes are generated in degree* 3*.*

Proof. Due to Lemma 3.1, it suffices to show that the bound holds for all cells. Let Z be a cell of an $(m \times n)$-transportation polytope and let $\mathcal{A} = Z \cap \mathbb{Z}^{m\times n}$. We may suppose that Z is full-dimensional, as all cells are faces of full-dimensional cells.

Let $d : \mathcal{A} \times \mathcal{A} \to \mathbb{N}$, $d(M, N) := |\{(i, j) \,|\, m_{ij} \neq n_{ij}\}|$ denote the Hamming distance of two matrices. For $\boldsymbol{u}, \boldsymbol{v} \in \mathbb{N}^{\mathcal{A}}$ with $\boldsymbol{x}^{\boldsymbol{u}} - \boldsymbol{x}^{\boldsymbol{v}} \in I_{\mathcal{A}}$, we define the distance $\bar{d}(\boldsymbol{u}, \boldsymbol{v}) := \min\{d(M, N) \,|\, M \in \operatorname{supp}(\boldsymbol{u}), N \in \operatorname{supp}(\boldsymbol{v})\}$.

Suppose that the theorem is wrong, *i. e.* there exists a binomial $\boldsymbol{x}^{\boldsymbol{u}} - \boldsymbol{x}^{\boldsymbol{v}} \in I_{\mathcal{A}}$ of degree $k \geq 4$ that cannot be expressed by binomials in $I_{\mathcal{A}}$ of smaller degree. Choose $\boldsymbol{u}$ and $\boldsymbol{v}$ s. t. $l := \bar{d}(\boldsymbol{u}, \boldsymbol{v})$ is minimal over all binomials with this property. Let $M^1 \in \operatorname{supp}(\boldsymbol{u})$ and $N^1 \in \operatorname{supp}(\boldsymbol{v})$ be matrices with $d(M^1, N^1) = l$. We can deduce $l \geq 4$ from the fact that row and column sums in M^1 and N^1 agree.

We will need the following lemma, which tells us something about the structure of M^1 and N^1:

Lemma 4.2. (i) There are no indices i_1, i_2, j_1, j_2 with $m^1_{i_1 j_1} = m^1_{i_2 j_2} = n^1_{i_1 j_2} = n^1_{i_2 j_1} = 1$ and $m^1_{i_1 j_2} = m^1_{i_2 j_1} = n^1_{i_1 j_1} = n^1_{i_2 j_2} = 0$, *i. e.* it is impossible, that at the same time

$$\begin{array}{c|cc} & j_1 & j_2 \\ \hline i_1 & 1 & 0 \\ i_2 & 0 & 1 \end{array} \text{ is a submatrix of } M^1 \text{ and } \begin{array}{c|cc} & j_1 & j_2 \\ \hline i_1 & 0 & 1 \\ i_2 & 1 & 0 \end{array} \text{ is a submatrix of } N^1. \quad (5)$$

(ii) This statement remains true, if we replace in one of the two submatrices a zero by a one or a one by a zero.

Proof. We will prove (i) and (ii) at the same time. Suppose that at least one of the statements is false. We will show that in this case, the distance of M^1 and N^1 is not minimal. Define a matrix $\tilde{M}$ by:

$$\tilde{m}_{ij} = \begin{cases} 1 & \text{for } (i,j) = (i_1, j_2) \text{ or } (i,j) = (i_2, j_1) \\ 0 & \text{for } (i,j) = (i_1, j_1) \text{ or } (i,j) = (i_2, j_2) \\ m_{ij} & \text{otherwise} \end{cases} \tag{6}$$

WLOG in case (ii), N^1 is the matrix where one entry was replaced. Hence, $\tilde{M} \in Z$. In both cases, the following inequality holds:

$$d(\tilde{M}, N^1) \leq d(M^1, N^1) - 2 \tag{7}$$

There exist matrices M^2 and M^3 in $\operatorname{supp}(\boldsymbol{u})$, s. t. for $A = M^2 + M^3$ the inequalities $a_{i_1 j_2} \geq \tilde{m}_{i_1 j_2} = 1$ and $a_{i_2 j_1} \geq \tilde{m}_{i_2 j_1} = 1$ hold. This follows from the fact that $\sum_{M \in \operatorname{supp}(\boldsymbol{u})} M = \sum_{N \in \operatorname{supp}(\boldsymbol{v})} N$.

It follows, that $(M^1 + M^2 + M^3 - \tilde{M})$ has only positive entries. Thus, it is contained in $2 \cdot Z$. By the Generalized Birkhoff-von Neumann Theorem (Theorem 2.1), there exist matrices A^2 and A^3 s. t. $M^1 + M^2 + M^3 = \tilde{M} + A^2 + A^3$.

Now choose $\boldsymbol{u'}$ s. t.

$$\boldsymbol{x}^{\boldsymbol{u'}} = \frac{x_{\tilde{M}} \cdot x_{A^2} \cdot x_{A^3}}{x_{M^1} \cdot x_{M^2} \cdot x_{M^3}} \boldsymbol{x}^{\boldsymbol{u}} \tag{8}$$

By construction, $\boldsymbol{x}^{\boldsymbol{u'}} - \boldsymbol{x}^{\boldsymbol{v}}$ is a binomial in $I_{\mathcal{A}}$ of degree k, that cannot be expressed by binomials of smaller degree. Using (7), we deduce $\bar{d}(\boldsymbol{u'}, \boldsymbol{v}) \leq d(\tilde{M}, N^1) < d(M^1, N^1) = \bar{d}(\boldsymbol{u}, \boldsymbol{v})$. Contradiction. □

Now we finish the proof of Theorem 4.1. As $M^1 \neq N^1$, there must be a position, in which M^1 has a one and N^1 has a zero. Since both matrices have the same column sums, there must be a position in the same column, where the situation is converse. So WLOG, we can write:

$$M^1 = \begin{bmatrix} 1 & \cdots & \cdots \\ 0 & \cdots & \cdots \\ & \cdots & \cdots \end{bmatrix} \qquad N^1 = \begin{bmatrix} 0 & \cdots & \cdots \\ 1 & \cdots & \cdots \\ & \cdots & \cdots \end{bmatrix} \tag{9}$$

As row sums in both matrices agree, there must be another entry in the first row, where N^1 has a one and M^1 has a zero. Likewise, there must be an entry in the second row, where N^1 has a zero and M^1 has a one. Using Lemma 4.2 we obtain that those entries must be in different columns. Thus, WLOG we can write:

$$M^1 = \begin{bmatrix} 1 & & 0 & \cdots \\ 0 & 1 & & \cdots \\ & \cdots & & \cdots \end{bmatrix} \qquad N^1 = \begin{bmatrix} 0 & & 1 & \cdots \\ 1 & 0 & & \cdots \\ & \cdots & & \cdots \end{bmatrix} \tag{10}$$

If $n^1_{23} = 0$ was valid, we would have a submatrix, which is forbidden by Lemma 4.2:

$$M^1 = \begin{bmatrix} 1 & & 0 & \cdots \\ 0 & 1 & & \cdots \\ & \cdots & & \cdots \end{bmatrix} \qquad N^1 = \begin{bmatrix} 0 & & 1 & \cdots \\ 1 & 0 & 0 & \cdots \\ & \cdots & & \cdots \end{bmatrix} \tag{11}$$

So, $n_{23} = 1$ must hold.

In the same way, we obtain the values for the other darkly shaded entries:

$$M^1 = \begin{bmatrix} 1 & 1 & 0 & \dots \\ 0 & 1 & 0 & \dots \\ & \dots & & \dots \end{bmatrix} \qquad N^1 = \begin{bmatrix} 0 & 0 & 1 & \dots \\ 1 & 0 & 1 & \dots \\ & \dots & & \dots \end{bmatrix} \tag{12}$$

Now, we can reapply this argument: due to the fact that the row sums in both matrices are the same, there is another entry in the first row where N^1 has a one and M^1 has a zero, as well as another entry in the second row in another column, where N^1 has a zero and M^1 has a one. Because of Lemma 4.2, the two entries above/below are fixed as well and we obtain:

$$M^1 = \begin{bmatrix} 1 & 1 & 0 & 1 & 0 & \dots \\ 0 & 1 & 0 & 1 & 0 & \dots \\ & & \dots & & & \dots \end{bmatrix} \qquad N^1 = \begin{bmatrix} 0 & 0 & 1 & 0 & 1 & \dots \\ 1 & 0 & 1 & 0 & 1 & \dots \\ & & \dots & & & \dots \end{bmatrix} \tag{13}$$

Reapplying this argument again and again we arrive at a point where all columns are “used up” and it is unavoidable that a submatrix that is forbidden by Lemma 4.2 appears. Contradiction. □

4.2. **Generalization to Flow Polytopes.**

In this subsection, we show that the Main Theorem can be deduced from the result on transportation polytopes proved in the previous subsection.

Proof of the Main Theorem. First note that Theorem 4.1 also holds for transportation polytopes that have integer lower and upper bounds on the entries of the matrices. This implies that the theorem holds for flow polytopes whose underlying graph is bipartite and all arcs point to the same color class.

Now let $F_G = F_{G,\boldsymbol{d},\boldsymbol{u},\boldsymbol{l}}$ be a flow polytope. We will see below that it is possible to transform an arbitrary graph $G = (V, A)$ into a bipartite graph $G' = (V', A')$, s. t. all arcs point to the same color class and there exists a bijective map $\varphi : F_G \cap \mathbb{Z}^A \to F_{G'} \cap \mathbb{Z}^{A'}$ which can be extendend to an isomorphism of the semigroups that those sets of lattice points generate. This implies, that the toric ideals I_{F_G} and $I_{F'_G}$ are equal. By the observation above, $I_{F_{G'}}$ is generated in degree 3.

The following transformation (essentially [11, 21.6a]) can be used: split every vertex $v \in V$ into two vertices v' and v''. For every arc (v, w), add an arc (v', w''), which inherits the upper and lower bounds of (v, w). For every $v \in V$, add arcs of the form (v', v''). Choose a constant $N \in \mathbb{N}$ that satisfies $N \geq \max\{\sum_{w:(v,w)\in V} u_{(v,w)}, \sum_{w:(w,v)\in V} u_{(w,v)}\}$ for all $v \in V$. Set $d'_{v'} = -N$, $d'_{v''} = N + d_v$ as well as $u'_{(v',v'')} = \infty$ and $l'_{(v',v'')} = 0$ for all $v \in V$. For an example see Figure 2. □

5. Gröbner Bases for Transportation Polytopes

In this section, we discuss degree bounds for Gröbner bases of transportation polytopes and we construct Gröbner bases in high degree.

For the toric ideal of the Birkhoff polytope, the following degree bound for their revlex Gröbner bases is known:

Theorem 5.1 ([2, Theorem 6.1],[12, Theorem 14.8])**.** *Let I_{B_n} be the toric ideal of the Birkhoff polytope B_n. Let $\mathcal{G}$ be a reduced Gröbner basis of I_{B_n} with respect to an arbitrary reverse lexicographic term order.*

Then, $\mathcal{G}$ has at most degree n.



(a) Before

(b) After

FIGURE 2. A graph $G = (V, A)$, its demand vector and a flow on the graph are transformed. For all $a \in A$, $l_a = 0$ and $u_a = 12$.

The proof of this theorem uses the fact that every integer matrix in B_n contains exactly n ones. Consider a cell Z of an $(m \times n)$-transportation polytope. Either for Z or for the complementary cell Z' obtained by exchanging zeroes and ones, the following statement holds: every integer matrix in the cell contains at most $\lfloor mn/2 \rfloor$ ones. Using this observation, essentially the same proof yields the following theorem:

Theorem 5.2. *Let $T_{\boldsymbol{rc}}$ be an $(m \times n)$-transportation polytope and let $\prec$ denote a term order that corresponds to a subdivide-and-pull triangulation (as defined in Section 3).*

Then, the reduced Gröbner basis of $I_{T_{\boldsymbol{rc}}}$ has at most degree $\left\lfloor \frac{m \cdot n}{2} \right\rfloor$.

This improves a known degree bound for reduced Gröbner bases ([12, Proposition 13.15]) by a factor of approximately 2.

Both theorems are (almost) as good as they can get, in the following sense:

Theorem 5.3 (B_n has revlex Gröbner bases in degree n)**.** *Fix $n \in \mathbb{N}$. Then there exists a revlex term order $\prec$, s. t. the reduced Gröbner basis $\mathcal{G}_\prec$ of $I_{B_{2n}}$ has exactly degree $2n$.*

Theorem 5.4 (Gröbner bases in high degree for transportation polytopes)**.** *Let m and n be even. Then there exists a smooth $(m \times n)$-transportation polytope $T_{\boldsymbol{rc}}$ and a term order $\prec$, s. t. the reduced Gröbner basis $\mathcal{G}_\prec$ of $I_{T_{\boldsymbol{rc}}}$ has degree at least $\frac{m(n-2)}{2}$.*

The term order can be chosen to be revlex or it can correspond to a subdivide-and-pull triangulation (as defined in Section 3).

Theorem 5.3 supports the experimental result of Diaconis and Eriksson, who suggested that revlex Gröbner bases of I_{B_n} have exactly degree n ([1, Remark 9]).

Examples for the constructions used in the proofs of Theorems 5.3 and 5.4 can be found at the end of this section.

Proof of Theorem 5.3. For $i = 1, \ldots, n$, define $(n \times n)$-matrices M^i, $\tilde{M}^i$, N^i and $\tilde{N}^i$ by $m^i_{ii} = 1$, $\tilde{m}^i_{i+1,i} = 1$ (here $n + 1 := 1$), $n^i_{jj} = 1$ for $j \neq i$ and all other entries are zero. $\tilde{N}_i$ is defined in the following way: column i and row $i + 1$ contain only zeroes. If this row and column are deleted, the remaining matrix is an identity matrix.

$$A_i = \begin{bmatrix} M^i & N^i \\ \tilde{N}^i & \tilde{M}^i \end{bmatrix} \qquad\qquad B_i = \begin{bmatrix} \tilde{M}^i & \tilde{N}^i \\ N^i & M^i \end{bmatrix} \tag{14}$$

By the Generalized Birkhoff-von Neumann Theorem (Theorem 2.1), there exist matrices $M_1, \dots, M_{n-1}$ s. t.

$$\sum_{i=1}^{n}(A^i + B^i) = I_{2n} + \sum_{j=1}^{n-1} M_j = \begin{bmatrix} J_n & S_n \\ S_n & J_n \end{bmatrix} \tag{15}$$

where

$$S_n = \begin{bmatrix} 2n-3 & 1 & & & & \\ 1 & 2n-4 & 1 & & & \\ & 1 & \ddots & \ddots & & \\ & & \ddots & 2n-4 & 1 & \\ & & & 1 & 2n-4 & 1 \\ & & & & 1 & 2n-3 \end{bmatrix} \tag{16}$$

$$J_n = \begin{bmatrix} 1 & & & & 1 \\ 1 & 1 & & & \\ & \ddots & \ddots & & \\ & & 1 & 1 & \\ & & & 1 & 1 \end{bmatrix} \tag{17}$$

and I_n denotes the identity matrix.

Order the lattice points of B_{2n} s. t. I_{2n} is the minimal element and the matrices A_i and B_i are smaller than all the remaining elements. In the revlex order $\prec$ defined by this ordering, the left side of equation (15) corresponds to the initial term. One can show that it is a minimal generator of the initial ideal $\mathrm{in}_{\prec}(I_{B_{2n}})$. This follows from two facts: each A_i and each B_i is necessary to cover all the ones in I_{2n} and each A_i and each B_i has a one that is not covered by any of the other A_is or B_is. □

Proof of Theorem 5.4. First, we define some notation for a couple of $\left(\frac{m}{2} \times \frac{n}{2}\right)$-matrices with $\{0,1\}$-entries. Let $\mathbf{1}$ and $\mathbf{0}$ denote the matrices where all entries are 1 or 0 respectively. Let $\mathbf{1_{ij}}$ be the matrix, whose entry at position (i,j) is 1 and all other entries are 0. Let $\mathbf{0_{ij}} := \mathbf{1} - \mathbf{1_{ij}}$.

Using those matrices, we will now define some $(m \times n)$-matrices:

$$A_{ij} = \begin{bmatrix} \mathbf{1_{i1}} & \mathbf{0_{ij}} \\ \mathbf{0_{i1}} & \mathbf{1_{ij}} \end{bmatrix} \qquad\qquad B_{ij} = \begin{bmatrix} \mathbf{1_{ij}} & \mathbf{0_{i1}} \\ \mathbf{0_{ij}} & \mathbf{1_{i1}} \end{bmatrix} \tag{18}$$

$$E = \begin{bmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{bmatrix} \qquad\qquad D = \begin{bmatrix} \mathbf{0} & \mathbf{1} \\ \mathbf{1} & \mathbf{0} \end{bmatrix} \tag{19}$$

$$C = \begin{bmatrix} 1 & 0 & \dots & 0 & 0 & 1 & \dots & 1 \\ \vdots & & \ddots & & \vdots & & \ddots & \\ 1 & 0 & \dots & 0 & 0 & 1 & \dots & 1 \\ 0 & 1 & \dots & 1 & 1 & 0 & \dots & 0 \\ \vdots & & \ddots & & \vdots & & \ddots & \\ 0 & 1 & \dots & 1 & 1 & 0 & \dots & 0 \end{bmatrix} \qquad \mathcal{B} := \{A_{ij}\} \cup \{B_{ij}\} \tag{20}$$

The following equation is a relation of lattice points in the polytope $T_{\boldsymbol{rc}}$ with $\boldsymbol{r} = \left(\frac{n}{2}, \dots, \frac{n}{2}\right)$ and $\boldsymbol{c} = \left(\frac{m}{2}, \dots, \frac{m}{2}\right)$:

$$\sum_{i=1}^{\frac{m}{2}} \sum_{j=2}^{\frac{n}{2}} (A_{ij} + B_{ij}) = \left(\frac{n}{2} - 2\right) \cdot C + \left(\frac{m(n-2)-n}{2} + 1\right) \cdot D + E = \tag{21}$$

$$\begin{bmatrix} \frac{n}{2} - 1 & 1 & \dots & 1 & \frac{m(n-2)-n}{2} + 1 & \frac{m(n-2)}{2} - 1 & \dots & \frac{m(n-2)}{2} - 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{n}{2} - 1 & 1 & \dots & 1 & \frac{m(n-2)-n}{2} + 1 & \frac{m(n-2)}{2} - 1 & \dots & \frac{m(n-2)}{2} - 1 \\ \frac{m(n-2)-n}{2} + 1 & \frac{m(n-2)}{2} - 1 & \dots & \frac{m(n-2)}{2} - 1 & \frac{n}{2} - 1 & 1 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{m(n-2)-n}{2} + 1 & \frac{m(n-2)}{2} - 1 & \dots & \frac{m(n-2)}{2} - 1 & \frac{n}{2} - 1 & 1 & \dots & 1 \end{bmatrix}$$

Let $\mathcal{A} = T_{\boldsymbol{rc}} \cap \mathbb{Z}^{m \times n}$. Choose a term order $\prec$ s. t. either

(i) $\prec$ is revlex and E is the minimal element and the elements of $\mathcal{B}$ are smaller than all the remaining elements of $\mathcal{A}$

(ii) or $\prec$ corresponds to a subdivide-and-pull triangulation, E is the minimal element in the full-dimensional cell Z that (21) is contained in and the elements of $\mathcal{B}$ are smaller than all the remaining elements of $Z \cap \mathcal{A}$.

The left side of (21) corresponds to the initial term in the order $\prec$. One can show that it is a minimal generator of the initial ideal $\mathrm{in}_{\prec}(I_{T_{\boldsymbol{rc}}})$. This follows from two facts: each element of $\mathcal{B}$ is necessary to cover all the ones in E and each element of $\mathcal{B}$ contains a one that is not covered by any other element of $\mathcal{B}$.

Hence, the reduced Gröbner basis of $I_{T_{\boldsymbol{rc}}}$ with respect to $\prec$ contains an element of degree $\frac{mn}{2}$.

Lemma 1 in [5] characterizes the marginals of smooth transportation polytopes. The transportation polytope defined above is not smooth. However, if we add mn to all entries in the last column of the matrices in (21), we obtain a relation of lattice points in the transportation polytope with marginals $\boldsymbol{r} = \left(\frac{n}{2} + mn, \dots, \frac{n}{2} + mn\right)$ and $\boldsymbol{c} = \left(\frac{m}{2}, \dots, \frac{m}{2}, m^2 n + \frac{m}{2}\right)$. This polytope is smooth. □

Example 5.5. This example shows that Theorem 5.3 holds for $n = 6$.

$$\underbrace{\begin{bmatrix} 1 & & & & & \\ & & & & 1 & \\ & & & & & 1 \\ & 1 & & & & \\ & & & 1 & & \\ & & 1 & & & \end{bmatrix}}_{A_1} + \underbrace{\begin{bmatrix} & & & 1 & & \\ & 1 & & & & \\ & & & & & 1 \\ 1 & & & & & \\ & & 1 & & & \\ & & & & 1 & \end{bmatrix}}_{A_2} + \underbrace{\begin{bmatrix} & & & 1 & & \\ & & & & 1 & \\ & & 1 & & & \\ & & & & & 1 \\ 1 & & & & & \\ & 1 & & & & \end{bmatrix}}_{A_3} + \underbrace{\begin{bmatrix} & & & & 1 & \\ 1 & & & & & \\ & & & & & 1 \\ & & & 1 & & \\ & 1 & & & & \\ & & 1 & & & \end{bmatrix}}_{B_1} + \underbrace{\begin{bmatrix} & & & 1 & & \\ & & & & & 1 \\ & 1 & & & & \\ 1 & & & & & \\ & & & & 1 & \\ & & 1 & & & \end{bmatrix}}_{B_2} + \underbrace{\begin{bmatrix} & & 1 & & & \\ & & & 1 & & \\ & & & & 1 & \\ 1 & & & & & \\ & 1 & & & & \\ & & & & & 1 \end{bmatrix}}_{B_3}$$

$$= \begin{bmatrix} 1 &&&&& \\ & 1 &&&& \\ && 1 &&& \\ &&& 1 && \\ &&&& 1 & \\ &&&&& 1 \end{bmatrix} + \begin{bmatrix} && 1 &&& \\ 1 &&&&& \\ & 1 &&&& \\ &&&&& 1 \\ &&& 1 && \\ &&&& 1 & \end{bmatrix} + 2 \cdot \begin{bmatrix} &&& 1 && \\ &&&& 1 & \\ &&&&& 1 \\ 1 &&&&& \\ & 1 &&&& \\ && 1 &&& \end{bmatrix} + \begin{bmatrix} &&&& 1 & \\ &&& 1 && \\ &&&&& 1 \\ 1 &&&&& \\ && 1 &&& \\ & 1 &&&& \end{bmatrix} + \begin{bmatrix} &&& 1 && \\ &&&&& 1 \\ &&&& 1 & \\ & 1 &&&& \\ 1 &&&&& \\ && 1 &&& \end{bmatrix} = \begin{bmatrix} 1 && 1 & 3 & 1 & \\ 1 & 1 && 1 & 2 & 1 \\ & 1 & 1 && 1 & 3 \\ 3 & 1 && 1 && 1 \\ 1 & 2 & 1 & 1 & 1 & \\ & 1 & 3 && 1 & 1 \end{bmatrix}$$

This equation corresponds to an element of I_{B_6}. We order the lattice points in B_6 s. t. I_6 is minimal and $A_1, A_2, A_3, B_1, B_2, B_3$ are smaller than all the remaining points. In the revlex order $\prec$ defined by this ordering, the left side of the equation corresponds to the initial term. One can show that it is a minimal generator of the initial ideal $\mathrm{in}_{\prec}(I_{B_6})$. Hence, the reduced Gröbner basis of I_{B_6} with respect to $\prec$ contains an element of degree 6.

Example 5.6. This example shows that Theorem 5.4 holds for $n = m = 6$. The following equation is a relation of lattice points in the polytope $T_{\boldsymbol{rc}}$ with $\boldsymbol{r} = \boldsymbol{c} = (3, 3, 3, 3, 3, 3)$. A translated version of this relation is contained in the transportation polytope with marginals $\boldsymbol{r} = (39, 39, 39, 39, 39, 39)$ and $\boldsymbol{c} = (3, 3, 3, 3, 3, 219)$. By [5, Lemma 1], this polytope is smooth.

$$\underbrace{\begin{bmatrix} 1 &&& 1 && 1 \\ &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ & 1 & 1 && 1 & \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \end{bmatrix}}_{A_{12}} + \underbrace{\begin{bmatrix} 1 &&& 1 & 1 & \\ &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ & 1 & 1 &&& 1 \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \end{bmatrix}}_{A_{13}} + \underbrace{\begin{bmatrix} &&& 1 & 1 & 1 \\ 1 &&& 1 && 1 \\ &&& 1 & 1 & 1 \\ 1 & 1 & 1 &&& \\ & 1 & 1 && 1 & \\ 1 & 1 & 1 &&& \end{bmatrix}}_{A_{22}} + \underbrace{\begin{bmatrix} &&& 1 & 1 & 1 \\ 1 &&& 1 & 1 & \\ &&& 1 & 1 & 1 \\ 1 & 1 & 1 &&& \\ & 1 & 1 &&& 1 \\ 1 & 1 & 1 &&& \end{bmatrix}}_{A_{23}} + \underbrace{\begin{bmatrix} &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ 1 &&& 1 && 1 \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \\ & 1 & 1 && 1 & \end{bmatrix}}_{A_{32}} + \underbrace{\begin{bmatrix} &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ 1 &&& 1 & 1 & \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \\ & 1 & 1 &&& 1 \end{bmatrix}}_{A_{33}}$$

$$+ \underbrace{\begin{bmatrix} & 1 &&& 1 & 1 \\ &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ 1 && 1 & 1 && \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \end{bmatrix}}_{B_{12}} + \underbrace{\begin{bmatrix} && 1 && 1 & 1 \\ &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ 1 & 1 && 1 && \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \end{bmatrix}}_{B_{13}} + \underbrace{\begin{bmatrix} &&& 1 & 1 & 1 \\ & 1 &&& 1 & 1 \\ &&& 1 & 1 & 1 \\ 1 & 1 & 1 &&& \\ 1 && 1 & 1 && \\ 1 & 1 & 1 &&& \end{bmatrix}}_{B_{22}} + \underbrace{\begin{bmatrix} &&& 1 & 1 & 1 \\ && 1 && 1 & 1 \\ &&& 1 & 1 & 1 \\ 1 & 1 & 1 &&& \\ 1 & 1 && 1 && \\ 1 & 1 & 1 &&& \end{bmatrix}}_{B_{23}} + \underbrace{\begin{bmatrix} &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ & 1 &&& 1 & 1 \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \\ 1 && 1 & 1 && \end{bmatrix}}_{B_{32}} + \underbrace{\begin{bmatrix} &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ && 1 && 1 & 1 \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \\ 1 & 1 && 1 && \end{bmatrix}}_{B_{33}}$$

$$= 1 \cdot \underbrace{\begin{bmatrix} 1 &&&& 1 & 1 \\ 1 &&&& 1 & 1 \\ 1 &&&& 1 & 1 \\ & 1 & 1 & 1 && \\ & 1 & 1 & 1 && \\ & 1 & 1 & 1 && \end{bmatrix}}_{C} + 10 \cdot \underbrace{\begin{bmatrix} &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \end{bmatrix}}_{D} + \underbrace{\begin{bmatrix} 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \\ 1 & 1 & 1 &&& \\ &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \\ &&& 1 & 1 & 1 \end{bmatrix}}_{E} = \begin{bmatrix} 2 & 1 & 1 & 10 & 11 & 11 \\ 2 & 1 & 1 & 10 & 11 & 11 \\ 2 & 1 & 1 & 10 & 11 & 11 \\ 10 & 11 & 11 & 2 & 1 & 1 \\ 10 & 11 & 11 & 2 & 1 & 1 \\ 10 & 11 & 11 & 2 & 1 & 1 \end{bmatrix}$$

We order the lattice points of $T_{\boldsymbol{rc}}$ s. t. E is minimal and the A_{ij}s and B_{ij}s are smaller than all the remaining points.

In the revlex order $\prec$ defined by this ordering, the left side of the equation corresponds to the initial term. One can show that it is a minimal generator of the initial ideal $\mathrm{in}_{\prec}(I_{T_{\boldsymbol{rc}}})$. Hence, the reduced Gröbner basis of $I_{T_{\boldsymbol{rc}}}$ with respect to $\prec$ contains an element of degree 12.

Acknowledgments

This article contains the main results of my *Diplomarbeit* (Master's thesis) [7], written in 2007 at Freie Universität Berlin. I would like to thank my advisor Christian Haase for many inspiring hints and discussions.

References

[1] Persi Diaconis and Nicholas Eriksson. Markov bases for noncommutative Fourier analysis of ranked data. *Journal of Symbolic Computation*, 41(2), February 2006. `arXiv:math/0405060v2`.

[2] Persi Diaconis and Bernd Sturmfels. Algebraic algorithms for sampling from conditional distributions. *Ann. Statist.*, 26(1):363–397, 1998. Available at `http://www-stat.stanford.edu/~cgates/PERSI/papers/sturm98.pdf`.

[3] Günter Ewald and Alexa Schmeinck. Representation of the Hirzebruch-Kleinschmidt varieties by quadrics. *Beiträge Algebra Geom.*, 34(2):151–156, 1993.

[4] Christian Haase. *Lattice Polytopes and Triangulations.* PhD thesis, Technische Universität Berlin, 2000. Available at `http://opus.kobv.de/tuberlin/frontdoor.php?source_opus=90`.

[5] Christian Haase and Andreas Paffenholz. Quadratic Gröbner bases for smooth 3 × 3 transportation polytopes. *Journal of Algebraic Combinatorics*, 30(4):477–489, 2009. `arXiv:math.CO/0607194`.

[6] Carl W. Lee. Regular triangulations of convex polytopes. In *Applied Geometry and Discrete Mathematics: The Victor Klee Festschrift*, volume 4 of *DIMACS Series in Discrete Math. and Theor. Comput. Sci.*, pages 443–456. American Mathematical Society, Providence, 1991.

[7] Matthias Lenz. Torische Ideale von Flusspolytopen. Master's thesis, Freie Universität Berlin, July 2007. `arXiv:0709.3570v3`.

[8] Hidefumi Ohsugi and Takayuki Hibi. Convex polytopes all of whose reverse lexicographic initial ideals are squarefree. *Proc. Amer. Math. Soc.*, 129(9):2541–2546 (electronic), 2001.

[9] J. Rambau, F. Santos Leal, and J. A. De Loera. *Triangulations: Structures for Algorithms and Applications*, volume 25 of *Algorithms and Computation in Mathematics.* Springer, 2010. Not yet published. Available: October 2010.

[10] Francisco Santos. MSRI, 1997.

[11] Alexander Schrijver. *Combinatorial Optimization – Polyhedra and Efficiency.* Number 24A in Algorithms and Combinatorics. Springer, Berlin, 2003.

[12] Bernd Sturmfels. *Gröbner Bases and Convex Polytopes*, volume 8 of *University Lecture Series.* American Mathematical Society, Providence, Rhode Island, 1996.

[13] Bernd Sturmfels. Equations defining toric varieties. Kollár, János (ed.) et al., Algebraic geometry. Proceedings of the Summer Research Institute, Santa Cruz, CA, USA, July 9–29, 1995. Providence, RI: American Mathematical Society. Proc. Symp. Pure Math. 62(pt.2), 437-449, 1997. `arXiv:alg-geom/9610018v1`.

[14] Seth Sullivant. Compressed polytopes and statistical disclosure limitation. *Tohoku Math. J. (2)*, 58(3):433–445, 2006. `arXiv:math.CO/0412535`.

[15] Günter M. Ziegler. *Lectures on Polytopes*, volume 152 of *GTM.* Springer-Verlag, New York, 1995.

Technische Universität Berlin, Sekretariat MA 4-5, Strasse des 17. Juni 136, 10623 Berlin
E-mail address: `lenz@math.tu-berlin.de`